

ARTERIAL HYPERTENSION AND THYROID STATUS IN PATIENTS OF DIFFERENT AGES

Khusainova Munira Alisherovna

Assistant of Samarkand State Medical University

Mavlonkulova Dilnura Mansurovna

Student of Samarkand State Medical University

Nizomova Dilnoza Erkinovna

Student of Samarkand State Medical University

Xusenova Ruxshona Farxodovna

Student of Samarkand State Medical University

Muhammadiyeva Malika Toxirovna

Student of Samarkand State Medical University

Rasulova Dildora Tolibovna

Student of Samarkand State Medical University

Erkinova Nafisa Yorkinovna

Student of Samarkand State Medical University

Samarkand State Medical University, Samarkand, Uzbekistan

ABSTRACT

174 patients of different genders and ages with arterial hypertension were examined. Among women under the age of 54, the incidence of combination of arterial hypertension with diffuse changes in the structure of the thyroid gland was higher than among men of the same age group (64.1% and 47.1%, respectively). Among women in The age group after 54 years saw an increase in the frequency of thyroid echostructure disorders to 72.5%, among men this figure was 42.4%. The incidence of nodular goiter among women under the age of 54 was 10.3%, after 54 years – 23.5%; among men - 5.9 and 6.1%, respectively. In the age groups up to 54 years, the frequency of hormonal signs of hypothyroidism among women with diffuse thyroid changes was 32.0 %, among men – 16.7%; in the groups after 54 years, these figures were 21.6 and 21.4%, respectively. A decrease in the functional activity of the thyroid gland in individuals with its diffuse changes led to an increase in the frequency and severity of cardiovascular pathology.

Key words: arterial hypertension, thyroid status, gender and age of patients

INTRODUCTION

In recent years, there has been a steady increase in the number of diseases of the cardiovascular system. First of all, this applies to arterial hypertension (AH) and its complications. Epidemiological studies have provided data confirming the effect of

hypertension on the incidence of atherosclerosis and associated coronary artery disease, the occurrence of coronary heart disease and heart failure. The main risk factors for hypertension include age, smoking, hypercholesterolemia and dyslipidemia, and a family history of early cardiovascular disease diseases, abdominal obesity, etc. Information has also been obtained on the aggravating role of disorders of the functional state of the pituitary-thyroid system in the development and progression AH. Parameters such as cardiac output, heart rate, and total peripheral vascular resistance depend to a certain extent on the function of the thyroid gland. A decrease in the level of thyroid hormones in the body contributes to the development of atherosclerosis, sodium and water retention in the body, accumulation of proteoglycans and glycoproteins, swelling of the vascular wall, impaired endothelial function, which eventually leads to an increase in blood pressure (BP). Against the background of latent, erased forms of thyroid pathology, in which the symptoms of endocrine disease may be unexpressed, and elevated blood pressure comes first, hypotensive therapy often has only short-term success due to uncorrected subclinical hypo- or hyperthyroidism (thyroid gland). With thyroid dysfunction in combination with hypertension, other concomitant somatic diseases, usually chronic, worsen or form. Considering the information that the frequency of pathology Thyroid gland among therapeutic patients is growing and has increased more than 9 times in recent years, the study of the structure and function of thyroid gland in patients with cardiovascular diseases is relevant. The purpose of this work is to study the features of the functional state of the pituitary-thyroid system and the echostructure of the thyroid gland in patients with arterial hypertension, depending on gender and age.

MATERIALS AND METHODS OF RESEARCH

174 patients of the therapeutic clinic with hypertension of II, III degrees were examined, among whom there were 90 women and 84 men aged 24 to 66 years. All the examined persons gave informed consent to participate in the study, which meets ethical standards, and the general condition of all patients upon admission was regarded as relatively satisfactory. Patients with arterial hypertension syndrome were not included in the study. hypertension, in which the increase in blood pressure was secondary, as well as patients with severe concomitant pathology (acute myocardial infarctions, heart defects, rhythm disturbances, bronchial asthma, diabetes mellitus). All patients received basic antihypertensive therapy from the first day of hospitalization. The median age of the examined individuals was 54 years, on the basis of which all patients – men and women - were divided by age into two groups: before 54 and after 54 years. Ultrasound of the heart revealed signs characterizing the severity of hypertension: hypertrophy of the left ventricle (LV), hypertrophy of the interventricular septum (MP), dilation of cavities, diastolic dysfunction. The determination of hormones of the pituitary-thyroid system in blood serum was carried

out by the radioimmune method using commercial kits: - thyroid-stimulating hormone (TSH): RIA TSH / CIS bio international, regulatory values 0.25-4.0 μ ed/ml; - thyroxine (T4): RIA-gnost T4 /CIC bio international, regulatory values 58-142 nmol/L; - free T4: RIA-gnost FT4 /CIC bio international, standard values 9.0-23.2 pmol/L; - triiodothyronine (T3): RIA-gnost T3 /CIC bio international, regulatory values 0.9-2.9 nmol/L; - free T3: RIA-gnost FT3 /CIC bio international, regulatory values 3.1-6.5 pmol/L.

THE RESULTS AND THEIR DISCUSSION

The ratio of women and men in the age groups of patients with hypertension before 54 years was 43.3 and 56.7%, and after 54 years – 60.7 and 39.3%, respectively ($p < 0.05$). Consequently, in the age group up to 54 years, hypertension was more often registered in men than in women, and in the age group after 54 years, on the contrary, it was detected more frequently in women than in men, which is consistent with the literature data. Among women under the age of 54, the incidence of combination of hypertension with diabetes was higher than in men of the same age group (64.1 and 47.1 % accordingly). After the age of 54, an increase in the frequency of thyroid echostructure disorders was noted among women to 72.5%; in men, this figure was 42.4%. In the age groups up to 54 years, in women with hypertension, the thyroid volume was 6.49 ± 0.15 ml, with hypertension and pancreas – 6.51 ± 0.25 ml, in men, the values of this indicator were 7.25 ± 0.12 and 7.31 ± 0.16 ml, respectively. In the age groups after 54 years, in women with hypertension, the volume of thyroid was 6.54 ± 0.16 ml, with hypertension and pancreas – 7.04 ± 0.2 ml, in men, the values of this indicator were 7.10 ± 0.13 and 7.27 ± 0.20 ml, respectively. There were no cases of excess of the thyroid gland volume above the standard values (more than 18 ml for women and 25 ml for men). On the contrary, rather low volume values attract attention Thyroid in the examined patients. In people with dysphoria in the age groups up to 54 years, among women, the ratio of frequencies of UNO and NO structure of thyroid was 32.0 and 68.0%, among men - 58.3 and 41.7%; in the age groups after 54 age among women - 40.5 and 59.5%, among men – 35.7 and 64.3%, respectively. The incidence of ultrasound among women under the age of 54 was 10.3 %, after 54 years – 23.5%; among men under the age of 54 – 5.9%, after 54 years – 6.1%.

Thus, the frequency of ultrasound increased by more than 2 times among women with increasing age, and the frequency of cases of thyroid disorders increased by 1.5 times among men with diabetes. Among the examined women with hypertension and diabetes in both age groups were those who took L-thyroxine (3 people in each group). Data on the serum levels of thyroid hormones and TSH in these women were not used to calculate the average values of these indicators for groups. The heterogeneous echostructure of the thyroid gland is a sign of tissue restructuring of the organ in response to pathological iodine deficiency or to the effects of others . strumogens (food

or ecopathogens). Inflammatory foci can also be visualized in the form of hypoechoic inclusions in the thyroid gland at the initial stages of the formation of the autoimmune process. At the same time, all intermediate types of thyroid echostructure that do not correspond to the description of a completely homogeneous structure of thyroid tissue, according to experts, are a visual reflection of various thyroid pathology. Received in the work The results indicate a high incidence of dysphoria among the examined patients with hypertension, not only women (68.9%), but also men (45.2%), which suggests a high prevalence of thyroid pathology among the population of Siberian territories, in which dysphoria, nodular forms of goiter and hypothyroidism occupy a significant place. What are the reasons for this phenomenon? As is known, one of the main reasons is iodine deficiency, but the data obtained in recent years indicate that as a result of government decisions At the level of measures for the prevention of iodine deficiency conditions, the provision of iodine to the Uzbek population has become more adequate. However, a rapid transition from iodine deficiency to increased iodine intake may adversely affect the structure and function of the thyroid gland. Structural changes in Thyroid can also be formed under the action of other specific and non- specific strumogens that damage the thyroid tissue and disrupt its function. Since disorders of the echostructure of thyroid tissue are a reflection of the emerging pathology of the thyroid gland, there is a high incidence among patients with dysphoria the frequency of hormonal signs of hypothyroidism, it can be argued that in patients with hypertension and hypertension, the risk of severe cardiovascular pathology is higher than in patients with hypertension in combination with normal thyroid structure and function.

CONCLUSIONS:

It has been shown that even hypothyroidism itself can cause hypertension, which in some patients is reversible against the background of levothyroxine replacement therapy. It is known that the prevalence of thyroid pathology among women is several times higher than among men. In the study, the ratio of the incidence of the among men and women with hypertension turned out to be 1 : 1.5. Consequently, the frequency of thyroid pathology among men is increasing, and this allows to say that they have an increased risk of developing a more severe course of diseases of the cardiovascular system due to an increase in the frequency of disorders of the structure and function of the thyroid gland.

LITERATURE

1. Khusainova, M. A., Eshmamatova, F. B., Ismoilova, K. T., & Mamadiyorova, M. M. (2023). METABOLIC SYNDROME IN RHEUMATOID ARTHRITIS AS A CRITERION OF CARDIOVASCULAR RISK. *Oriental renaissance: Innovative, educational, natural and social sciences*, 3(1), 331-339.

2. Khusainova, M. A. (2023). CYSTATIN C IS AN EARLY MARKER OF DECREASED KIDNEY FUNCTION. *Oriental renaissance: Innovative, educational, natural and social sciences*, 3(1), 485-490.
3. Khusainova, M. A., Vakhidov, J. J., Khayitov, S. M., & Mamadiyorova, M. M. (2023). Cardiac arrhythmias in patients with rheumatoid arthritis. *Science and Education*, 4(2), 130-137.
4. Khusainova, M. A. (2023). Comorbidity thyrotoxicosis with coronary heart disease. *Science and Education*, 4(5), 205-213.
5. Khusainova, M. A., Gafforov, X. X., Eshmamatova, F. B., & Khayitov, S. M. (2023). Assessment of the quality of life in patients with exogenous allergic alveolitis. *Science and Education*, 4(2), 145-152.
6. Khusainova, M. A., Khaydarov, S. N., Makhmudova, K. D., & Ortikova, S. X. (2023). Features of prevention of chronic kidney diseases and chronic heart failure. *Science and Education*, 4(5), 242-250.
7. Khusainova, M. A., Khaydarov, S. N., Makhmudova, K. D., & Nayimov, A. S. (2023). Prevalence of bronchiolitis in patients with Rheumatoid arthritis. *Science and Education*, 4(5), 232-241.
8. Yarmatov, S., Khusainova, M., & Djabbarova, N. (2023). STUDY OF QUALITY OF LIFE INDICATORS IN PATIENTS WITH CORONARY HEART DISEASE USING THE SF-36 QUESTIONNAIRE. *Бюллетень студентов нового Узбекистана*, 1(7), 58-64.
9. Alisherovna, K. M., Totlibayevich, Y. S., Xudoyberdiyevich, G. X., & Jamshedovna, K. D. (2022). CLINICAL FEATURES OF HEART FAILURE IN PATIENTS WITH ISCHEMIC HEART DISEASE AND THYROTOXICOSIS. *Spectrum Journal of Innovation, Reforms and Development*, 7, 108-115.
10. Alisherovna, K. M., & Xudoyberdiyevich, G. X. Features of Heart Damage in Patients with Viral Cirrhosis of the Liver. *International Journal of Innovations in Engineering Research and Technology*, 8(04), 53-57.
11. Alisherovna, K. M., Totlibayevich, Y. S., Xudoyberdiyevich, G. X., & Jamshedovna, K. D. (2022). EFFICACY OF DRUG-FREE THERAPY OF HYPERTENSION DISEASES IN THE EARLY STAGE OF THE DISEASE. *Spectrum Journal of Innovation, Reforms and Development*, 7, 82-88.
12. Alisherovna, K. M., Toshtemirovna, E. M. M., & Oybekovna, E. E. (2022). Quality of life of patients with cirrhosis of the liver. *Spectrum Journal of Innovation, Reforms and Development*, 4, 197-202.
13. Alisherovna, K. M., Kairatovna, R. A., Umirovna, I. S., & Oybekovich, T. M. (2023). CHRONIC OBSTRUCTIVE PULMONARY DISEASE AND

- ANEMIA. *Spectrum Journal of Innovation, Reforms and Development*, 21, 140-147.
14. Khabibovna, Y. S., & Alisherovna, K. M. (2024). STRESS TESTING IN PATIENTS WITH CORONARY HEART DISEASE. *Journal of new century innovations*, 45(3), 28-33.
 15. Alisherovna, K. M., Kulmuxammatovich, Y. U., Boymamatovna, E. F., & Shokirovich, S. A. (2023). THE STATE OF NEUROPEPTIDE-CYTOKINE STATUS IN ISCHEMIC HEART DISEASE. *Spectrum Journal of Innovation, Reforms and Development*, 11, 42-50.
 16. Alisherovna, K. M., Nizamitdinovich, K. S., Rustamovich, T. D., & Haqnazarovich, K. S. (2022). Mental Status and Quality of Life in Patients With Sinus Node Weakness Syndrome and Chronic Coronary Heart Failure of Ischemic Etiology. *Texas Journal of Medical Science*, 15, 78-82.
 17. Alisherovna, K. M., Erkinovna, K. Z., Djamshedovna, K. D., & Nizamitdinovich, K. S. (2023). QUALITY OF LIFE PATIENTS WITH OSTEOARTHRITIS. *Journal of new century innovations*, 36(1), 164-175.
 18. Khabibovna, Y. S., Alisherovna, K. M., Nizamitdinovich, K. S., Tashtemirovna, E. M. M., Abdukadirovna, A. S., & Jasurovna, J. S. (2023). DEPRESSION, ANXIETY AND QUALITY OF LIFE IN PATIENTS WITH ATRIAL FIBRILLATION. *Journal of new century innovations*, 39(1), 185-189.
 19. Khabibovna, Y. S., Alisherovna, K. M., Tashtemirovna, E. M. M., Totlibayevich, Y. S., Nizamitdinovich, K. S., & Baxtiyorovich, U. J. (2023). DIAGNOSTIC VALUE OF CYSTATIN C IN PATIENTS WITH HYPERTENSION AND OBESITY. *World Bulletin of Public Health*, 22, 55-59.
 20. Alisherovna, K. M., Khabibovna, Y. S., Nizamitdinovich, K. S., & Bakhtiyorovich, U. J. (2023). CYSTATIN and KIDNEY FUNCTION. *Journal of new century innovations*, 38(2), 220-225.
 21. Alisherovna, K. M., Djamshedovna, K. D., Totlibayevich, Y. S., & Boymamatovna, E. F. (2022). The Effectiveness of the Original Drug Trimetazidine MV in Patients with Stable Ischemic Heart Disease and Persistent Angina Attacks Against the Background of the Use of Trimetazidine Generics. *Miasto Przyszłości*, 30, 235-238.
 22. Khabibovna, Y. S., Alisherovna, K. M., Tashtemirovna, E. M. M., Nizamitdinovich, K. S., & Abdukadirovna, A. S. (2023). ANTITHROMBOTIC THERAPY IN CARDIOLOGICAL PATIENTS. *Journal of new century innovations*, 39(1), 169-171.
 23. Alisherovna, K. M., & Djamshedovna, K. D. (2024). AFTER COVID-19 QUALITY OF LIFE. *Spectrum Journal of Innovation, Reforms and Development*, 25, 103-110.

24. Alisherovna, K. M., Erkinovna, K. Z., Davranovna, M. K., & Pulotovna, Z. D. (2022). Positive Effect of Sorbitol in Patients with Chronic Renal Insufficiency. *Miasto Przyszłości*, 30, 214-217.
25. Alisherovna, K. M., Akmalovna, K. N., & Mamasoliyevna, D. N. (2022). Kidney dysfunction in chronic heart failure. *Texas Journal of Medical Science*, 13, 104-109.
26. Nizamitdinovich, K. S., Khabibovna, Y. S., Alisherovna, K. M., & Tashtemirovna, E. M. M. (2023). Spinal Injury for Rheumatoid Arthritis. *Miasto Przyszłości*, 40, 426-432.
27. Baxtiyorovich, U. J., Alisherovna, K. M., & Mamasoliyevna, D. N. (2023). Features of cognitive impairment in patients with chronic kidney disease at predialysis stages. *World Bulletin of Public Health*, 22, 49-54.
28. Khabibovna, Y. S., Alisherovna, K. M., Erkinovna, K. Z., & Djamshedovna, K. D. (2023). Gender Characteristics of the Course of Rheumatoid Arthritis. *Miasto Przyszłości*, 40, 438-442.
29. Erkinovna, K. Z., Alisherovna, K. M., Bakhtiyorovich, U. J., & Djamshedovna, K. D. (2023). METABOLIC SYNDROME IN RHEUMATOID ARTHRITIS. *Journal of new century innovations*, 38(2), 203-211.
30. Alisherovna, K. M., Totlibayevich, Y. S., Xudoyberdiyevich, G. X., & Jamshedovna, K. D. (2022). PSYCHOSOMATIC FEATURES AND THE LEVEL OF DEPRESSION WITH CHRONIC HEART FAILURE IN PATIENTS WITH ARTERIAL HYPERTENSION AND CORONARY HEART DISEASE. *Spectrum Journal of Innovation, Reforms and Development*, 7, 89-95.
31. Alisherovna, K. M., Baxtiyorovich, Z. M., & Anvarovich, N. J. (2022). To Assess The Condition Of The Myocardium In Patients Chronic Heart Failure On The Background Of Rheumatoid Arthritis. *Spectrum Journal of Innovation, Reforms and Development*, 4, 210-215.
32. Alisherovna, K. M., Toshtemirovna, E. M. M., Djamshedovna, K. D., & Maxammadiyevich, K. S. (2022). Endothelial Dysfunction in Patients with Chronic Heart Failure. *Miasto Przyszłości*, 30, 218-220.
33. Tashtemirovna, E. M. M., Khabibovna, Y. S., Alisherovna, K. M., & Erkinovna, K. Z. (2023). Angiopathy in Rheumatoid Arthritis. *Miasto Przyszłości*, 40, 418-425.
34. Alisherovna, K. M., Akramovna, I. K., & Yorkinovna, E. N. (2024). CLINICAL AND MORPHOLOGICAL CRITERIA OF COLITIS IN PATIENTS WITH CHRONIC ISCHEMIC DISEASE OF THE DIGESTIVE SYSTEM. *Ta'lim innovatsiyasi va integratsiyasi*, 18(6), 6-13.
35. Alisherovna, K. M., Akramovna, I. K., & Kairatovna, R. A. (2024). THE EFFECTIVENESS OF TREATMENT OF PATIENTS WITH OSTEOARTHRITIS WITH CARDIOVASCULAR DISORDERS IN

- METABOLIC SYNDROME. *Ta'lim innovatsiyasi va integratsiyasi*, 18(5), 223-230.
36. Akramovna, I. K., & Alisherovna, K. M. (2024). CAUSES OF ARRHYTHMIA DURING PREGNANCY. *Journal of new century innovations*, 45(3), 34-41.
37. Mamasoliyevna, D. N., Akmalovna, K. N., & Alisherovna, K. M. (2022). Quality of Life Depending on Gender. *The Peerian Journal*, 11, 71-77.
38. Mamasoliyevna, D. N., Alisherovna, K. M., & Totlibayevich, Y. S. (2023). Diabetes Mellitus and Non-Alcoholic Fatty Liver Disease: the Facets of Conjugacy. *Miasto Przyszłości*, 35, 166-173.
39. Alisherovna, K. M., Akramovna, I. K., & Baxtiyorovna, O. K. (2024). THE COURSE OF CHRONIC ISCHEMIC PANCREATITIS IN PATIENTS WITH CORONARY HEART DISEASE. *Ta'lim innovatsiyasi va integratsiyasi*, 18(5), 231-239.
40. Alisherovna, K. M., & Yusupovich, N. F. (2021). FEATURES OF THE CIRCADIAN RHYTHM BLOOD PRESSURE IN PATIENTS HEART FAILURE AND IMPAIRED RENAL FUNCTION. *EDITOR COORDINATOR*, 423.
41. Alisherovna, K. M., Totlibayevich, Y. S., Davranovna, M. K., & Erkinovna, K. Z. (2022). ASSESSMENT OF THE GENERAL CONDITION AND QUALITY OF LIFE OF WOMEN POST-PRODUCTIVE AGE, WHO SUFFERED FROM COVID-19 ASYMPTOMATICALLY, AND 12 MONTHS AFTER THE MODERATE SEVERE FORM OF THE DISEASE. *Spectrum Journal of Innovation, Reforms and Development*, 10, 277-282.
42. Alisherovna, K. M., Sherzodovna, M. D., Tursunboyevna, I. K., & Uktamovna, U. U. (2023). LEFT VENTRICULAR HYPERTROPHY IN PERSONS WITHOUT ARTERIAL HYPERTENSION: PSYCHOSOMATIC APPROACH TO THE STUDY OF THIS PHENOMENON.
43. Alisherovna, M. K. Stages Of Development Of Renal Dysfunction And Anemia In Patients With Chronic Heart Failure. *International Journal of Innovations in Engineering Research and Technology*, 8(05), 50-53.
44. Alisherovna, M. K., & Xudoyberdiyevich, G. X. (2021). Treatment of Chronic Heart Diseases Insufficiency Depending On the Condition of the Kidneys. *International Journal of Innovations in Engineering Research and Technology*, 8(05), 64-69.